

Summary of Overexpressor G482, Family CAAT

Mendel Biotechnology, Inc.

[Summary](#) | [Sequence](#) | [Expression](#) | [Morphology](#) | [Physiology](#) | [Biochemistry](#) | [Microarray](#)

Published Information

- [Executive Summary](#)
 - [Overview](#)
 - [November 2000 Notes](#)
- Traits**

- [November 2000 Traits](#)
- [August 2000 Traits](#)
- [April 2000 Traits](#)
- [November 1999 Traits](#)
- [All Traits](#)
- [Corrigenda](#)

Genes

- [November 2000 Genes](#)
- [Index by Gene ID](#)
- [Index by Family](#)
- [Index by Keyword](#)
- [DNA FASTA files](#)

Assays

- [Gene Expression](#)
- [Morphology](#)
- [Physiology](#)
- [Biochemistry](#)
- [Microarrays](#)

Approach

- [Gene Determination](#)
- [Overexpression](#)
- [Knockouts](#)
- [Vector Information](#)
- [Bioinformatics](#)
- [Growth Facilities](#)

People

- [Staff](#)
- [Advisory Board](#)
- [Home Page](#)

Mendel Discoveries

The function of G482 was analyzed at Mendel through its ectopic overexpression in plants. G482 overexpressors are more tolerant to high NaCl in a germination assay. Further experiments should be done to determine the utility of this degree of NaCl tolerance under field conditions.

RT-PCR analysis of endogenous levels of G482 transcripts indicate that this gene is expressed constitutively in all tissues tested. A cDNA array experiment supports the RT-PCR derived tissue distribution data. G482 is not induced above basal levels in response to any environmental stress treatments tested.

Closely Related Genes from Other Species

The closest homology in the non-Arabidopsis plant database is within the conserved domain of G482 and therefore no potentially orthologous genes are available in the public domain.

Utilities

The potential utilities of this gene include the ability to confer salt tolerance during the germination stage of a crop plant. This would most likely impact survivability and yield. Evaporation of water from the soil surface causes upward water movement and salt accumulation in the upper soil layer, where the seeds are placed. Thus, germination normally takes place at a salt concentration much higher than the mean salt concentration in the whole soil profile.

Gene and Trait Disclosure Chronology

November 1999 Analyzed as Overexpressor

November 1999 Increased tolerance to high salt

References

Edwards D, et al. Multiple genes encoding the conserved CCAAT-box transcription factor complex are expressed in Arabidopsis. *Plant Physiol.* 1998 Jul;117(3):1015-22.

Keywords

Salt, Germination

Knockout Status

KO DNA insertion not identified

Plasmid ID P47

Cloning Vector pMEN20

Cloning Site SalINotI

Source DNA N97233

Bacterial Strain DH5a

Scientist	Jacqueline Heard
Contact	Phone: (510) 264-0280 x112 Fax: (510) 264-0254 Email:
Information	jheard@mendelbio.com

Patent Information

Plant Trait Modification I, 11/17/99

Polynucleotides for Seed Trait Alteration, 3/22/00

Fri Nov 10 18:52:47 PST 2000

The information provided in this data release is proprietary and confidential to Mendel Biotechnology, Inc.